













Erratum

Chuc NT, Larsson M, Do NT, Diwan VK, Tomson GB, Falkenberg T. Improving private pharmacy practice: A multi-intervention experiment in Hanoi, Vietnam. *J Clin Epidemiol* 2002;55(11):1148–55.

Table 1 in the aforementioned article was published with missing elements. The corrected table is presented below.

doi: 10.1016/S0895-4356(03)00173-2

Table 1: Mean (%) and standard deviation (SD) of encounters for ARI, STD, Cefalexin and Prednisolone tracer variables, Summary statistics and test of difference between the intervention (I) and control (C) groups across the study time and Trend-line graphs with the means (%) plotted out.

Variables		Mean % (SD)				Summary statistics		Trend-line graphs (%)	
		BL	RE	ED	PI	Mean*10 ⁻² (S.E*10 ⁻³)	p-value 2-tailed		
A	Antibiotics dispensed	I	45 (50)	26 (44)	15 (36)	30 (46)	-5.6 (2.3)	0.023	
		C	39 (49)	30 (46)	30 (46)	42 (50)	1.0 (1.8)		
R	Questions related to breathing	I	11 (31)	6 (23)	24 (43)	30 (46)	6.7 (2.6)	0.006	
		C	10 (30)	7 (26)	8 (27)	7 (25)	-1.4 (0.9)		
I	Acceptable Drugs Paracetamol, TM & cough syrups	I	44 (50)	50 (50)	61 (49)	48 (50)	0.6 (14.0)	0.731	
		C	33 (47)	38 (49)	35 (48)	29 (46)	-1.0 (12.2)		
S	Advised the client to go to a doctor	I	22 (42)	1 (10)	38 (49)	37 (49)	10 (3.0)	0.01	
		C	20 (40)	3 (16)	15 (36)	18 (38)	0.7 (1.4)		
T	Question regarding sexual activity	I	22 (24)	16 (26)	12 (17)	30 (25)	7.1 (49.3)	0.32	
		C	24 (24)	21 (24)	15 (21)	37 (22)	23.4 (59.4)		
D	Advised condom use	I	1 (12)	1 (10)	6 (25)	7 (24)	2.2 (1.1)	0.46	
		C	1 (9)	0 (0)	1 (9)	4 (21)	1.1 (0.4)		
	Correct symptomatic treatment	I	3(17)	7(26)	21(41)	30(46)	17.6 (2.9)	0.01	
		C	4(19)	9(29)	15(36)	19(39)	6.4 (1.9)		
C E F A L E X I N	Cefalexin dispensed	I	95 (23)	91 (29)	60 (49)	56 (50)	6.7 (1.9)	0.002	
		C	94 (23)	89 (31)	79 (41)	89 (31)	1.1 (0.7)		
	Prescription request	I	0 (0)	6 (23)	13 (33)	21 (41)	6.7 (1.9)	0.009	
		C	0 (0)	0 (0)	5 (22)	2 (15)	1.1(0.7)		
P R E D N I S O L O N E	Prescription request	I	1 (12)	2 (14)	20 (40)	18 (39)	6.7 (1.5)	0.002	
		C	1 (8)	1 (12)	4 (20)	4 (21)	1.3 (0.8)		
	Steroids dispensed	I	78 (42)	65 (48)	29 (46)	17 (37)	-22.0 (2.2)	0.001	
		C	73 (44)	69 (46)	64 (48)	58 (50)	-5.1 (2.3)		
	No drugs, NSAID or TM dispensed	I	20(22)	33(27)	66(39)	83 (26)	21.0 (12.3)	0.001	
		C	26(29)	27(29)	31(31)	44 (35)	5.0 (11.3)		

BL: Baseline; RE: Regulatory enforcement; ED: Education; PI: Peer influence. Summary statistics mean: Mean of trends over all pharmacies in (I) or (C) group and p-value for difference in trend between the groups. (-): Trends of intervention group; (--) Trend of control group; NSAID: Non Steroid Anti Inflammatory Drugs; TM: Traditional Medicines